
AD-A193 383

MOLECULAR BIOLOGY OF ARCHAEBACTERIA(U) BRITISH COLUMBIA 1/1
UNIV VANCOUVER DEPT OF BIOCHEMISTRY P P DENNIS
31 MAR 88 N00014-87-G-0049

F/G 6/2

NL



AD-A193 383		16. RESTRICTIVE			FILE CUP	
N 2b. DECLASSIFICATION / DOWNGRADING SCHEDU	3. DISTRIBUTION AVAILABILITY OF REPORT  DISTRIBUTION STATEMENT X					
NA NA		Approved for public released  Distribution Unlimited and  S. MONITORING ORGANIZATION REPORT NUMBER(S)				
4. PERFORMING ORGANIZATION REPORT NUMBER	R(S)	5. MONITORING	<del>ORGANIZATION</del>	-REPORT	NUMBER(S)	
NA NA						
Univ. of British Columbia	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION ELECTER			7	
Oc. ADDRESS (City, State, and ZIP Code) Department of Biochemistry 2146 Health Sciences Mall Vancouver, B.C. V6T 1W5		7b. ADDRESS (Ci	ty, State, and ZI	PCo	MAR 2 9 1988	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMEN		DENTIFI	CATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code)	ONR	N00014-87-G-0049  10. SOURCE OF FUNDING NUMBERS				
800 N. Quincy St. Arlington, VA 22217-5000		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UN ACCESSION	
11. TITLE (Include Security Classification)		61153N	RR04106			
Molecular Biology of Archaeb  12. PERSONAL AUTHOR(S)  Patrick  13a. TYPE OF REPORT  13b. TIME C	P. Dennis	Lia pare or orne			he are come	
Annual FROM 4		14. DATE OF REPORT (Year, Month, Day) 15. P 880331		15. PAGE COUNT		
16. SUPPLEMENTARY NOTATION  17. COSATI CODES  FIELD GROUP SUB-GROUP	18. SUBJECT TERMS	(Continue on reven	se if necessary a	nd iden	tify by block number)	
19. ABSTRACT (Continue on reverse if necessary	and identify by block	number)		<del></del>	···	
Genomic restriction frag Sulfolobus acidocaldorius (Sa Ll2e were cloned and sequence frames identified. The deduc with eubacterial and eucaryot evolutionary relationships we of the purified superoxide di oligonucleotide was prepared sequence has been aligned to been characterized and the re radicals has been demonstrate	ed. The sequence of L10e and L12 ic proteins and re defined. The smutase enzymes and the gene wa other SOD seque gulation of trained and character	es were analy e protein section their structer N-terminal from Hou was cloned and notes. The gonscription (inited.	in Llle, Ll yzed and th quences wer ture-functi amino acid s determine sequenced. ene transcr	e, L10 e read e alignon and seque d, an The ipts l	0e and ding gned d ence nave gen	
20. ÖRTRIBEHENPÄVAIDABEITF OF GAM FRORT		5				
22a. NAME OF RESPONSIBLE INDIVIDUAL		IZZD. IELEPHONE	(include Area Co	de)   22c	. OFFICE SYMBOL	

All other editions are obsolete.

Abbreviations: Hcu - Halobacterium cutirubrum

Hha - Halobacterium holobium (same species as Hcu)

Hvo - Halobacterium volcanii Sac - Sulfolobus acidocaldorius

Eco - Escherichia coli

#### Research Objectives

(i) to characterize the principles of gene organization and regulation of gene expression in archaebacteria; (ii) to elucidate the evolutionary relationship between these novel organisms and the traditional eubacterial and eucaryotic organisms; (iii) to understand in biophysical and molecular terms some of the mechanisms that allow archaebacteria to inhabit extreme environments.

# Progress - Year 1

Ribosomal protein genes: The ribosoml A protein complex forms the stalk structure on the large ribosome subunit and is comprised of four copies (two dimers) of L12e and one copy of L10e ribosomal protein. The L10e protein either binds to ribosomal RNA either directly or through the L11e protein which forms a bulge at the base of the stalk. This domain on the large subunit is the site of factor binding and associated GTPase activities during the protein synthesis cycle and is a conserved and defined feature on the ribosome from all organisms (1). In Eco the genes encoding the L11, L10 and L12 ribosomal proteins are located within a 3.0 Kb region of genomic DNA. This region has been cloned, sequenced and extensively characterized (2,3).

Partial amino acid sequences of many archaebacterial ribosomal proteins have been determined (4). We used the sequences from the Hcu L11e and L12e proteins to deduce and synthesize oligonucleotide sequences complementary to the corresponding genes. We cloned initially a 1.2 Kb Pst-Bam fragment encoding L12e and subsequently a 5.2 Kb Cla-Bam fragment containing both L11e and L12e. The sequence of the entire 5.2 Kb fragment was determined and it was found to contain several long open reading frames. Four of these open reading frames were shown to encode the L11e, L10e and L12e ribosomal proteins. The gene order as present in Eco is conserved in Hcu although the transcriptional organization will certainly be much different (work in progress).

Using a similar strategy the corresponding genomic region of Sac has been cloned and sequenced. It contains the same four genes in the same order. From the Hcu and Sac nucleic acid sequence we have deduced the amino acid sequences for the four ribosoml proteins and have begun to align the protein to their eubacterial and eucaryotic equivalent and deduce their evolutionary relationships. This analysis has been completed for the L10e and L12e protein (5).

There are presently four complete sequences of L10e proteins (one eubacterial, two archaebacterial and one eucaryotic) and 16 complete sequences of L12e (eight eubacterial, three archaebacterial and five eucaryotic). Using these sequences we have carried out inter-kingdom L10e and L12e alignments. The L10e protein from the three kingdoms were found to be co-linear. (A cDNA sequence published by Rich and Steitz was shown by us to encode the human L10e

protein (6)). The eubacterial protein is much shorter than the archaebacterial and eucaryotic proteins because of two large deletions, one internal and one at the carboxy terminus. The two archaebacterial L10e proteins were most similar; they exhibit 27% amino acid identity with only a single deletion-insertion. Inter-kingdom comparisons exhibit 15-25% amino acid identity with 5-7 deletion insertions.

STATE OF THE PROPERTY OF THE P

の後のでは、「これ」というできる。

The L12e protein from archaebacteria and eucaryotes can also be aligned and are co-linear. The eubacterial protein could not be made to fit this pattern although two regions appear to have homologous domain in the archaebacterial and eucaryotic proteins. The C-terminal domain of the Eco L12e protein has been shown to contain (i) a conserved face for interaction with extrinsic translation factors; (ii) an anion (potential GTP) binding site; and (iii) a putative Ll2e-Ll2e dimerization site (1). This domain is separated from the N-terminus by an unstructured ala-pro-rich hinge sequence and aligns to a region near the amino terminus of the archaebacterial and eucaryotic proteins. Inter-kingdom comparisons range from 18-36 percent amino acid sequence identity over their domain. The second region conserved in all proteins is the ala-pro-rich sequence. In eubacteria this sequence precedes the facter interaction domain; in archaebacteria and eucaryotic it follows the facter interaction domain. The N-terminus of the eubacterial L12e protein exhibits more similarity to its own carboxy end than to any region within the archaebacterial-eucaryotic protein. The carboxy termini of the archaebacterial and eucaryotic protein are highly similar to each other and apparently not represented in the eubacterial L12e protein.

Intraspecies comparisons between L10e and L12e proteins indicate that the L10e protein of archaebacteria and eucaryotes contains a partial copy of the L12e protein fused to its carboxy terminus. (In eubacteria most of this fusion has been removed from L10e by the carboxy terminal deletion and the L12e protein has been restructured.) For the two Sac proteins the 31 carboxy teminal residues of L10e and L12e are identical; conservation at the nucleotide level is also perfect. This homology extends further into the central regions of the L10e and L12e proteins and includes a 26 amino acid long module, reiterated thrice in archaebacterial L10e, twice in eucaryotic L10e and once in the corresponding L12e protein. Careful examination indicates that a single copy of the modular sequence is also present in the L12e and L10e protein of eubacteria. This modular sequence may play a role in L12e dimerization, L10e-L12e complex formation and the function of the L10e-L2e complex in translation.

Based upon these alignments and shared homologies a model has been constructed to depict the evolution of the primordial L10e and L12e genes and proteins in the three phylogenetic groups (5).

Superoxide dismutase: Previously we had purified and characterized the SOD enzyme activity from Hcu (7). An N-terminal amino acid sequence was determined, an oligonucleotide was synthesized and the gene encoding the SOD enzyme was cloned as a 1.1 Kb Sau3A fragment and sequenced. The fragment contain a single long open reading frame encoding a protein of 200 amino acids. The first 56 codons of the ORF correspond to the amino acid sequence determined from the purified protein. The gene encodes a typical Mn/Fe ion type enzyme as evidenced by the fact that (i) it has an acidic pI; (ii) it







2 Codes

Jud til



contains no cys residues; (iii) it is relatively abundant in tyr and trp residues; (iv) it exhibits 41% amino acid identity with the Mn SOD of Bacillus stearothermophilus and (v) it conserves the four residues, his 29, his 76, asp 158 and his 162 that in the Bacillus enzyme (position 26, 81, 163 and 167 respectively) are utilized for Mn binding.

By S1 nuclease protection and primer extension analysis we have shown that the SOD mRNA is about 242 nucleotides in length. It begins two nucleotides in front of the ATG translation initiation codon and terminates about 40 nucleotides downstream of the TAA translation termination codon, in a T5 sequence that is preceded by a GC-rich sequence. This terminator sequence appears similar to other Hcu terminators. Analysis of this and other 5' flanking sequences have yet to reveal the concensus sequence involved in promoter recognition by RNA polymerase in Hcu (that is, there is little conservation between the bacterio-opsin, SOD, rRNA and r-protein 5' flanking sequences).

Like bacterio-opsin, the mRNA encoding SOD starts only two nucleotides in front of the ATG translation initiation codon. There is no homolog of the Shine Dalgarno purine-rich sequence of eubacteria that is complementary to the 3° end of 16S rRNA on the SOD mRNA. In contrast some r-protein genes of Hcu retain this sequence. Thus it seems that Hcu ribosomes can initiate translation by two separate mechanisms: at the first AUG on the transcript (SOD and bacterio-opsin) or by a Shine Dalgarno interaction at internal AUG codons.

We have discovered that the SOD gene in Hcu is subject to regulation. Addition of paraquat, a generator of oxygen radicals, to an exponential phase culture results in the 5-10x induction of SOD enzyme activity. After several days of growth in the presence of paraquat the SOD activity returns to normal. The culture has become resistant to the inhibitor and the frequency suggests that resistance arises by transposition insertion (as in bacterio-opsin negative mutants of Hha; 8) into a gene specifying sensitivity (i.e. transport of paraquat across the cell envelope). This observation is being investigated.

#### References

- (1) Liljas, A. (1982) Prog. Biophys. Mol. Biol. 40, 161-228.
- (2) Post, L., Strycharz, G., Nomura, M., Lewis, H. and Dennis, P. (1979) PMAS 76, 1697-1701.
- (3) Downing, W. and Dennis, P. (1987) J. Mol. Biol. 194, 609-620.
- (4) Matheson, A. (1985) The Bacteria 8, 761-777.
- (5) Shimmin, L., Ramirez, C., Hatheson, A. and Dennis, P. (1988) submitted for publication.
- (6) Rich, B. and Steitz, J. (1988) Mol. Cell Biol., in press.
- (7) May, B. and Dennis, P. (1987) J. Bacteriol. 169, 1417-1422.
- (8) Betlach, M., Leong, D. and Boyer, H. (1986) System. Appl. Microbiol. 7, 83-89.

# Objectives for Year 2

### Gene encoding r-protein

- 1) Carry out inter-kingdom alignment and analysis of Lile and Lie protein;
- 2) Define the transcriptional organization of the L11e, L1e, L10e and L12e gene clusters in Hcu and Sac, map 5' and 3' transcript ends, cap 5' transcript ends and analyze for conserved promoter and terminator sequences.

### Superoxide dismutase

- Cap mRNA and characterize induction by paraquat at the level of transcription;
- 4) Look for DNA binding protein to the SOD promoter region by in vivo foot-printing;
- 5) Clone and characterize SOD gene from a genetically stable strain Hvo, isolate paraquat-sensitive (null) and resistant (constitutive) mutants;
- 6) Utilize the transformation system of Hvo to characterize the genetic regulation of the SOD gene.

# Processing of rRNA

- 7) Characterize promoter and processing signals in the separate 16S-23S and 5S transcription units of Sac;
- 8) Utilize cloned copies of the 16S inverted repeat processing signal from Hcu to develop an assay for RNaseIII-like activity;
- 9) Attempt to purify Hcu RNaseIII-like activity, clone the gene and study substrate specificity by <u>in vitro</u> site-specific mutagenesis of cloned 16S inverted repeat sequence.

### **Publications**

May, B. and P. Dennis (1987) Superoxide dismutase from the extremely halophilic archaebacterium <u>Halobacterium cutirubrum</u>. J. Bacteriol. 169, 1417-1422.

Shimmin, L., C. Ramirez, A. Matheson and P. Dennis (1988) Sequence alignment and evolutionary comparison of the L10e and L2e ribosomal proteins from archaebacteria, eubacteria and eucaryotics. Submitted for publication (copy enclosed to ONR).

#### Training

Bruce May: MRC, predoctoral fellow, Canadian

Lawrence Shimmin: ONR, predoctoral student, Canadian

Craig Newton: ONR technician, Canadian

Diedre de Jong-Wong: ONR technician, Canadian

Phelgun Joshi: MRC predoctoral student (grant), Canadian Peter Durovic: MRC predoctoral student (grant), Canadian

### MOLECULAR BIOLOGY PROGRAM DISTRIBTUION L

ANNUAL, FINAL, AND TECHNICAL REPORTS (On Dr. C. E. Ballou Department of Biochemistry University of California Berkeley, CA 94720

Dr. Rita Colwell Department of Microbiology University of Maryland College Park, MD 20742

Dr. Lacy Daniels Department of Microbiology University of Iowa Iowa City, IA 52242

Dr. Patrick P. Dennis Department of Biochemistry University of British Columbia 2146 Health Sciences Mall Vancouver, B.C. V6T 1W5

Dr. Henryk Eisenberg The Weizmann Institute of Science Department of Polymer Research P.O. Box 26 Rehovot 76100, Israel

Dr. James G. Ferry
Department of Anaerobic Microbiology
Virginia Polytechnic Institute
and State University
Blacksburg, Virginia 24061

Dr. Robert P. Gunsalus Department of Microbiology UCLA 405 Hilgard Avenue Los Angeles, CA 90024

Dr. Ramesh Gupta Southern Illinois University Department of Chemistry and Biochemistry Carbondale, IL 62901

Dr. Holger W. Jannasch Woods Hole Oceanographic Institution Woods Hole, MA 02543

Dr. Jordan Konisky University of Illinois 809 South Wright Street Champaign, IL 61820

Dr. Mary E. Lidstrom Center for Great Lakes Studies University of Wisconsin-Milwaukee Milwaukee, WI 53201

Dr. Ralph Mitchell Division of Applied Sciences Harvard University 125 Pierce Hall Cambridge, MA 02138

Dr. Leonard Muscatine Department of Biology University of California Los Angeles, California 90024

Dr. David P. Nagle Department of Botany and Microbiology University of Oklahoma Norman, OK 73019 Dr. Gary J. Olsen Department of Biology Indiana University Jordan Hall 138 Bloomington, Indiana 47405

Dr. Gregory J. Olson Inorganic Materials Division Center for Materials Science National Bureau of Standards Washington, DC 20234

Dr. Leo Parks Department of Microbiology North Carolina State University Raleigh, NC 27695

Dr. John N. Reeve Department of Microbiology Ohio State University 484 West 12th Avenue Columbia, OH 43210-1292

Dr. V. Romanovsky
Office d'Etudes Marines et Atmospherique
64 rue Gabrielle
1180, Bruxelles, Belgium

Dr. Saul Roseman Department of Biology Johns Hopkins University Baltimore, MD 21218

Dr. David C. White Department of Microbiology Institute of Applied Microbiology University of Tennessee Knoxville, TN 37996-0845

Dr. Carl R. Woese Genetics Department University of Illinois 505 S. Goodwin Avenue Urbana, IL 61801

Dr. Ralph S. Wolfe 131 Burrill Hall University of Illinois Urbana, IL 61801

Dr. Eli D. Schmell, Code 1141MB Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000

Dr. Michael T. Marron, Code 1141MB Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000

Dr. Margo G. Haygood Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000

Adminstrator (2 copies, Enclose DTIC Form 50) Defense Technical Information Center Building 5, Cameron Station Alexandria, VA 22314

ANNUAL AND FINAL REPORTS ONLY (One copy each)
Commander
Chemical and Biological Sciences Division
Army Research Office
P. O. Box 12211
Research Triangle Park, NC 27709

Directorate of Life Sciences Air Force Office of Scientific Research Bolling Air Force Base Washington, DC 20332

Chemistry and Atmospheric Sciences Directorate Air Force Office of Scientific Research Bolling Air Force Base Washington, DC 20332

Director Biotechnology Division CRDEC Aberdeen Proving Grounds, MD 21010

Administrative Contracting Officer
ONR Resident Representative
(Address varies - obtain from your business office)

Director, Code 12 Applied Research and Technology Directorate Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000

Director, Code 22 Support Technology Directorate Office of Naval Technology 800 North Quincy Street Arlington, VA 22217-5000

Director, Code 112 Environmental Sciences Directorate Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000

Director, Code 113 Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000

HALLE SECTION OF THE SECTION OF THE

FINAL AND TECHNICAL REPORTS ONLY
Director (6 copies)
Naval Research Laboratory
Attn: Technical Information Division, Code 2627
Washington, DC 20375